

Supplemental Figures

Figure S1




Figure S1A. wt SARS-CoV-2 neutralizing antibody ID₈₀ titers measured by pseudovirus viral neutralizing antibody (PsVNA); shown and summarized by age group as in Figure 1A. Samples with titers below the level of detection were assigned a value of 10. **Figure S1B:** PsVNA ID₈₀ titers by household, using the sequence and color scheme shown for Figure 1B. Statistical differences in antibody titers were compared using the Mann-Whitney-Wilcoxon test (two-sided). A Bonferroni correction was applied to adjust for 3 age group comparisons and the p-value was multiplied by 2. One specimen from a child in the 0-4 years age group was not tested for neutralizing antibodies due to insufficient sample volume.

Figure S2




Figure S2A: PsVNA ID₅₀ titers **Figure S2B:** PsVNA ID₈₀ titers. Data are shown and summarized by age group as in Figure 1A. Statistical differences in antibody titers were compared using the Mann-Whitney-Wilcoxon test (two-sided). A Bonferroni correction was applied to adjust for 3 age group comparisons and the p-value was multiplied by 2. One specimen from a child in the 0-4 year age group was not tested for neutralizing antibodies due to insufficient sample volume.

SEARCH Study Team*

*Walker Black³, Christine Council-DiBitetto¹, Tina Ghasri¹, Amanda Gormley¹, Milena Gatto¹, Maria Jordan¹, Karen Loehr¹, Jason Morsell¹, Jennifer Oliva¹, Jocelyn San Mateo¹, Khadija Smith¹, Kimberli Wanionek¹, Cathleen Weadon¹, Suzanne Woods¹

Affiliations

1. Department of International Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, MD
2. Vaccine Research Center, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD
3. Centers for Disease Control and Prevention, Atlanta, GA